

SOCIAL ANXIETY SYMPTOMS ASSESSMENT QUESTIONNAIRE – RCADS-SP

Version for children and adolescents aged 8 to 17 years old

Patient's last name			File number			
Patient's f	irst name					
Health ins	Health insurance number				Year	Month
			E	хр.		
Date of	Year	Month	D	ay	Sex	
birth					M	F
Address (no., street)				ПХ	
City			Postal Code			
School						
Grade *						

How often do each of these things happen to you?

- 1. Answer each item based on the last month or the period of time since your last appointment.
- 2. Use the scale at the top of the table.
- 3. Answer each item by checking the box that represents your situation the best.

	Never	Sometimes	Often	Always
Items	0	1	2	3
 I worry when I think I have done poorly at something. 	□ 0	<u> </u>	_ 2	□ 3
2. I feel scared when I have to take a test.	□ 0	<u> </u>	_ 2	□ 3
3. I feel worried when I think someone is angry with me.	<u> </u>	<u> </u>	<u> </u>	□ 3
4. I worry that I will do badly at my school work.	□ 0	<u> </u>	_ 2	□ 3
5. I worry I might look foolish.	□ 0	<u> </u>	_ 2	□ 3
6. I worry about making mistakes.	□ 0	<u> </u>	_ 2	□ 3
7. I worry what other people think of me.	□ 0	<u> </u>	_ 2	□ 3
8. I feel afraid if I have to talk in front of my class.	<u> </u>	<u> </u>	<u> </u>	□ 3
I feel afraid that I will make a fool of myself in front of people.	□ 0	<u> </u>	□ 2	□ 3

Revised Children's Anxiety and Depression Scale - Social Phobia Subscale - RCADS-SP © 2003 Bruce F. Chorpita

Questionnaire completed by:			Date :		
Signature	Year	Month	Day		

^{* 3}rd grade of elementary school to 1st year of Cegep or college

Patient's last name	Patient's first name	File number

Section reserved for the practitioner	
Total raw score	
Total raw score	
Total number of items	9
Number of answered items (≥ 7)*	
Adjusted score	
Score T **	
Is the T score greater than the clinical cut-off value of 65?	☐ Yes ☐ No
Practitioner's analysis and commentary:	

Questionnaire reviewed by:			Date	Date:		
Practitioner's last name	Practitioner's first name	Licence number	Signature	Year	Month	Day

^{*} If 3 or more answers are missing, the score of the subscale cannot be used.

^{**} For children and adolescents aged 8 to 17 years old, use the conversion table to identify the T score of the subscale according to the patient's sex, school grade and adjusted raw score. Only the raw score can be used for patients aged 18 years and over.