

GENERALIZED ANXIETY SYMPTOMS ASSESSMENT QUESTIONNAIRE – RCADS-P-GAD

Version for parents or caregivers of children and adolescents aged 3 to 17 years old

| - | | | | | |
|---------------------|------------|-------------|-----|-----------|-------|
| Patient's last name | | File number | | | |
| Patient's first ı | name | | | | |
| Health insurar | nce number | Е | xp. | Year | Month |
| Date of birth | Year Month | D | ay | Sex M | F |
| Address (no., | street) | | | □ X | |
| City | | | | Postal Co | ode |
| | • | | | | |
| Caregiver | Last name | | F | irst name | |

How often do each of these things happen to your child?

1. Answer each item based on the last month or the period of time since your child's last appointment.

School Grade *

- 2. Use the scale at the top of the table.
- 3. Answer each item by checking the box that represents your child's situation the best.

| | Never | Sometimes | Often | Always |
|---|----------|-----------|-------|--------|
| Items | 0 | 1 | 2 | 3 |
| 1. My child worries about things. | □ 0 | <u> </u> | _ 2 | □ 3 |
| My child worries that something awful will happen to someone in the family. | <u> </u> | <u> </u> | □ 2 | □ 3 |
| My child worries that bad things will happen to him (her). | <u> </u> | <u> </u> | _ 2 | □ 3 |
| My child worries that something bad will happen to him (her). | □ 0 | <u> </u> | _ 2 | □ 3 |
| My child worries about what is going to happen. | □ 0 | 1 | _ 2 | □ 3 |
| 6. My child thinks about death. | □ 0 | <u> </u> | □ 2 | □ 3 |

Revised Children's Anxiety and Depression Scale – Generalized Anxiety Disorder Subscale - RCADS-P-GAD © 2003 Bruce F. Chorpita

| Questionnaire completed by: | | Date : | |
|-----------------------------|------|--------|-----|
| Signature | Year | Month | Day |

^{* 3}rd grade of elementary school to 1st year of Cegep or college

| Patient's last name | Patient's first name | File number |
|---------------------|----------------------|-------------|
| | | |

| Section reserved for the practitioner | · |
|---|------------|
| Total raw score | |
| Total number of items | 6 |
| Number of answered items (≥ 4)* | |
| Adjusted score = | |
| Score T ** | |
| Is the T score greater than the clinical cut-off value of 65? | ☐ Yes ☐ No |
| Practitioner's analysis and commentary: | |
| | |
| | |
| | |
| | |
| | |
| | |

| Questionnaire reviewed by: | | | Date: | Date: | | |
|----------------------------|---------------------------|----------------|-----------|-------|-------|-----|
| Practitioner's last name | Practitioner's first name | Licence number | Signature | Year | Month | Day |

^{*} If 3 or more answers are missing, the score of the subscale cannot be used.

^{**} For parents of children and adolescents aged 8 to 17 years old, use the conversion table to identify the T score of the subscale according to the patient's sex and school grade, in addition to the parent's raw score. Only the raw score can be used for children aged between 3 to 7 years old and patients aged 18 years and over.